

Riemann-Lagrange geometry for starfish/coral dynamical system

Mircea Neagu

Abstract

In this paper we develop the Riemann-Lagrange geometry, in the sense of nonlinear connection, d-torsions, d-curvatures and jet Yang-Mills entity, associated with the dynamical system concerning social interaction in colonial organisms.

Mathematics Subject Classification (2010): 53C43, 53C07, 83C50.

Key words and phrases: 1-jet spaces, jet least squares Lagrangian functions, jet Riemann-Lagrange geometry, starfish/coral dynamics.

1 Social interactions in colonial organisms

Let $m \geq 2$ be an integer. We introduce social interactions for starfish/coral dynamics as follows [1]:

$$\left\{ \begin{array}{l} \frac{dN^1}{dt} = \lambda_1 N^1 - \alpha_1 (N^1)^2 - \alpha_2 \left(\frac{m}{m-1} \right) \cdot N^1 N^2 + \\ \quad + \frac{\alpha_1}{m-1} \left(\frac{N^2}{N^1} \right)^{m-2} \cdot (N^2)^2 - \delta_1 F N^1 \\ \frac{dN^2}{dt} = \lambda_2 N^2 - \alpha_2 (N^2)^2 - \alpha_1 \left(\frac{m}{m-1} \right) \cdot N^1 N^2 + \\ \quad + \frac{\alpha_2}{m-1} \left(\frac{N^1}{N^2} \right)^{m-2} \cdot (N^1)^2 - \delta_2 F N^2 \\ \frac{dF}{dt} = \beta F (N^1 + N^2) + \gamma F^2 - \rho F, \end{array} \right. \quad (1)$$

where

- $\alpha_1, \alpha_2, \lambda_1, \lambda_2, \delta_1, \delta_2, \beta, \gamma, \rho$ are positive coefficients;
- N^1, N^2 are coral densities;
- F is the starfish density;
- λ_1 and λ_2 are growth rates;

- λ_1/α_1 and λ_2/α_2 are single species carrying capacities;
- β , δ_1 and δ_2 are the interaction coefficients for starfish preying on corals;
- γ is the coefficient of starfish aggregation.

Note that m is the effect of increasing the social parameter. If we set $m = 2$, we obtain the (2 corals/1 starfish)-model of Antonelli and Kazarinoff [2], in which every term of degree greater than one is quadratic. It is $m \geq 3$ which forces the social interaction terms to be nonquadratic.

The dynamical system (1) can be extended to a dynamical system of order two coming from a first order Lagrangian of least square type. This extension is called in the literature of specialty as the geometric dynamical system (see Udriște [7]).

2 The Riemann-Lagrange geometry

The system (1) can be regarded on the 1-jet space $J^1(\mathbb{R}, \mathbb{R}^3)$, whose coordinates are

$$\left(t, x^1 = N^1, x^2 = N^2, x_3 = F, y_1^1 = \frac{dN^1}{dt}, y_1^2 = \frac{dN^2}{dt}, y_1^3 = \frac{dF}{dt} \right).$$

Remark 1 We recall that the transformations of coordinates on the 1-jet space $J^1(\mathbb{R}, \mathbb{R}^3)$ are given by

$$\begin{cases} \tilde{t} = \tilde{t}(t) \\ \tilde{x}^i = \tilde{x}^i(x^j) \\ \tilde{y}_1^i = \frac{\partial \tilde{x}^i}{\partial x^j} \frac{dt}{dt} \cdot y_1^j, \end{cases} \quad (2)$$

where $i, j = \overline{1, 3}$.

In this context, the solutions of class C^2 of the system (1) are the global minimum points of the jet least square Lagrangian [3], [6]

$$\begin{aligned} L = & \left(y_1^1 - X_{(1)}^{(1)}(N^1, N^2, F) \right)^2 + \left(y_1^2 - X_{(1)}^{(2)}(N^1, N^2, F) \right)^2 + \\ & + \left(y_1^3 - X_{(1)}^{(3)}(N^1, N^2, F) \right)^2, \end{aligned} \quad (3)$$

where

$$\begin{aligned} X_{(1)}^{(1)}(N^1, N^2, F) &= \lambda_1 N^1 - \alpha_1 (N^1)^2 - \alpha_2 \left(\frac{m}{m-1} \right) \cdot N^1 N^2 + \\ &+ \frac{\alpha_1}{m-1} \left(\frac{N^2}{N^1} \right)^{m-2} \cdot (N^2)^2 - \delta_1 F N^1 \end{aligned}$$

$$\begin{aligned}
X_{(1)}^{(2)}(N^1, N^2, F) &= \lambda_2 N^2 - \alpha_2 (N^2)^2 - \alpha_1 \left(\frac{m}{m-1} \right) \cdot N^1 N^2 + \\
&\quad + \frac{\alpha_2}{m-1} \left(\frac{N^1}{N^2} \right)^{m-2} \cdot (N^1)^2 - \delta_2 F N^2 \\
X_{(1)}^{(3)}(N^1, N^2, F) &= \beta F (N^1 + N^2) + \gamma F^2 - \rho F,
\end{aligned}$$

Remark 2 *The solutions of class C^2 of the system (1) are solutions of the Euler-Lagrange equations attached to the the jet least square Lagrangian (3), namely (geometric dynamics)*

$$\frac{\partial L}{\partial x^i} - \frac{d}{dt} \left(\frac{\partial L}{\partial y_1^i} \right) = 0, \quad \forall i = \overline{1, 3}.$$

But, the jet least square Lagrangian (3) provides us with an entire Riemann-Lagrange geometry on the 1-jet space $J^1(\mathbb{R}, \mathbb{R}^2)$, in the sense of nonlinear connection, d-torsions, d-curvatures and jet Yang-Mills entity. Let us use the notation

$$J(X_{(1)}) = \left(\frac{\partial X_{(1)}^{(i)}}{\partial x^j} \right)_{i,j=\overline{1,3}} = \begin{pmatrix} J_{11} & J_{12} & J_{13} \\ J_{21} & J_{22} & J_{23} \\ J_{31} & J_{32} & J_{33} \end{pmatrix},$$

where

$$\begin{aligned}
J_{11} &= \lambda_1 - 2\alpha_1 N^1 - \alpha_2 \left(\frac{m}{m-1} \right) \cdot N^2 - \alpha_1 \left(\frac{m-2}{m-1} \right) \frac{(N^2)^m}{(N^1)^{m-1}} - \delta_1 F, \\
J_{12} &= -\alpha_2 \left(\frac{m}{m-1} \right) \cdot N^1 + \alpha_1 \left(\frac{m}{m-1} \right) \frac{(N^2)^{m-1}}{(N^1)^{m-2}}, \\
J_{13} &= -\delta_1 N^1, \\
J_{21} &= -\alpha_1 \left(\frac{m}{m-1} \right) \cdot N^2 + \alpha_2 \left(\frac{m}{m-1} \right) \frac{(N^1)^{m-1}}{(N^2)^{m-2}}, \\
J_{22} &= \lambda_2 - 2\alpha_2 N^2 - \alpha_1 \left(\frac{m}{m-1} \right) \cdot N^1 - \alpha_2 \left(\frac{m-2}{m-1} \right) \frac{(N^1)^m}{(N^2)^{m-1}} - \delta_2 F, \\
J_{23} &= -\delta_2 N^2, \\
J_{31} &= \beta F, \\
J_{32} &= \beta F, \\
J_{33} &= \beta (N^1 + N^2) + 2\gamma F - \rho.
\end{aligned}$$

Following the geometrical ideas from Balan and Neagu [3], we obtain the following geometrical results:

Theorem 3 (i) *The canonical nonlinear connection on $J^1(\mathbb{R}, \mathbb{R}^3)$, produced by the system (1), has the local components*

$$\Gamma = \left(M_{(1)1}^{(i)} = 0, N_{(1)j}^{(i)} \right),$$

where $N_{(1)j}^{(i)}$ are the entries of the skew-symmetric matrix

$$N_{(1)} = -\frac{1}{2} [J(X_{(1)}) - {}^T J(X_{(1)})] = \begin{pmatrix} N_{(1)1}^{(1)} & N_{(1)2}^{(1)} & N_{(1)3}^{(1)} \\ N_{(1)1}^{(2)} & N_{(1)2}^{(2)} & N_{(1)3}^{(2)} \\ N_{(1)1}^{(3)} & N_{(1)2}^{(3)} & N_{(1)3}^{(3)} \end{pmatrix},$$

where

$$N_{(1)1}^{(1)} = N_{(1)2}^{(2)} = N_{(1)3}^{(3)} = 0,$$

$$N_{(1)2}^{(1)} = -N_{(1)1}^{(2)} = \left(\frac{m}{m-1} \right) (\alpha_1 N^2 - \alpha_2 N^1) + \left(\frac{m}{m-1} \right) \left[\alpha_2 \frac{(N^1)^{m-1}}{(N^2)^{m-2}} - \alpha_1 \frac{(N^2)^{m-1}}{(N^1)^{m-2}} \right],$$

$$N_{(1)3}^{(1)} = -N_{(1)1}^{(3)} = \beta F + \delta_1 N^1,$$

$$N_{(1)3}^{(2)} = -N_{(1)2}^{(3)} = \beta F + \delta_2 N^2.$$

(ii) *All adapted components of the canonical generalized Cartan connection CT , produced by the system (1), are zero.*

(iii) *The effective adapted components $R_{(1)jk}^{(i)}$ of the torsion d-tensor $\mathbf{T}$ of the canonical generalized Cartan connection CT , produced by the system (1), are the entries of the following skew-symmetric matrices:*

$$R_{(1)1} = \left(R_{(1)j1}^{(i)} \right)_{i,j=\overline{1,3}} = \frac{\partial N_{(1)}}{\partial N^1} = \begin{pmatrix} 0 & \frac{\partial N_{(1)2}^{(1)}}{\partial N^1} & \delta_1 \\ -\frac{\partial N_{(1)2}^{(1)}}{\partial N^1} & 0 & 0 \\ -\delta_1 & 0 & 0 \end{pmatrix},$$

where

$$\frac{\partial N_{(1)2}^{(1)}}{\partial N^1} = \left(\frac{m}{m-1} \right) \left[-\alpha_2 + \alpha_2 (m-1) \left(\frac{N^1}{N^2} \right)^{m-2} + \alpha_1 (m-2) \left(\frac{N^2}{N^1} \right)^{m-1} \right];$$

$$R_{(1)2} = \left(R_{(1)j2}^{(i)} \right)_{i,j=\overline{1,3}} = \frac{\partial N_{(1)}}{\partial N^2} = \begin{pmatrix} 0 & \frac{\partial N_{(1)2}^{(1)}}{\partial N^2} & 0 \\ -\frac{\partial N_{(1)2}^{(1)}}{\partial N^2} & 0 & \delta_2 \\ 0 & -\delta_2 & 0 \end{pmatrix},$$

where

$$\frac{\partial N_{(1)2}^{(1)}}{\partial N^2} = \left(\frac{m}{m-1} \right) \left[\alpha_1 - \alpha_2 (m-2) \left(\frac{N^1}{N^2} \right)^{m-1} - \alpha_1 (m-1) \left(\frac{N^2}{N^1} \right)^{m-2} \right];$$

$$R_{(1)3} = \left(R_{(1)j3}^{(i)} \right)_{i,j=\overline{1,3}} = \frac{\partial N_{(1)}}{\partial F} = \begin{pmatrix} 0 & 0 & \beta \\ 0 & 0 & \beta \\ -\beta & -\beta & 0 \end{pmatrix}.$$

(iv) All adapted components of the curvature d -tensor $\mathbf{R}$ of the canonical generalized Cartan connection $C\Gamma$, produced by the system (1), cancel.

(v) The geometric electromagnetic-like distinguished 2-form, produced by the system (1), is given by

$$\mathbb{F} = F_{(i)j}^{(1)} \delta y_1^i \wedge dx^j,$$

where

$$\delta y_1^i = dy_1^i - N_{(1)j}^{(i)} dx^j, \quad \forall i = \overline{1,3},$$

and the adapted components $F_{(i)j}^{(1)}$ are the entries of the skew-symmetric matrix

$$F^{(1)} = \left(F_{(i)j}^{(1)} \right)_{i,j=\overline{1,3}} = -N_{(1)}.$$

(vi) The jet geometric Yang-Mills entity, produced by the system (1), is given by the formula

$$\begin{aligned} \mathcal{EYM}(t) &= \left[F_{(1)2}^{(1)} \right]^2 + \left[F_{(1)3}^{(1)} \right]^2 + \left[F_{(2)3}^{(1)} \right]^2 = \\ &= \left\{ \left(\frac{m}{m-1} \right) (\alpha_1 N^2 - \alpha_2 N^1) + \left(\frac{m}{m-1} \right) \left[\alpha_2 \frac{(N^1)^{m-1}}{(N^2)^{m-2}} - \alpha_1 \frac{(N^2)^{m-1}}{(N^1)^{m-2}} \right] \right\}^2 + \\ &\quad + (\beta F + \delta_1 N^1)^2 + (\beta F + \delta_2 N^2)^2. \end{aligned}$$

Remark 4 We recall that, under a transformation of coordinates (2), the local components of the nonlinear connection obey the rules (see [3], [4], [5])

$$\begin{aligned} \widetilde{M}_{(1)1}^{(k)} &= M_{(1)1}^{(j)} \left(\frac{dt}{\widetilde{dt}} \right)^2 \frac{\partial \widetilde{x}^k}{\partial x^j} - \frac{dt}{\widetilde{dt}} \frac{\partial \widetilde{y}_1^k}{\partial t}, \\ \widetilde{N}_{(1)l}^{(k)} &= N_{(1)i}^{(j)} \frac{dt}{\widetilde{dt}} \frac{\partial x^i}{\partial \widetilde{x}^l} \frac{\partial \widetilde{x}^k}{\partial x^j} - \frac{\partial x^i}{\partial \widetilde{x}^l} \frac{\partial \widetilde{y}_1^k}{\partial x^i}. \end{aligned}$$

Open problem. There exist real meanings in trophodynamics for the geometrical objects constructed in this paper?

References

- [1] P. Antonelli, P. Auger, R. Bradbury, *Corals and starfish waves on the Great Barrier Reef: Analytical trophodynamics and 2-patch aggregation methods*, Mathl. Comput. Modelling, Vol. **27**, No. **4** (1998), 121-135.
- [2] P. Antonelli, N. Kazarinoff, *Starfish predation of a growing coral reef community*, J. Theor. Biol. **107** (1984), 667-684.
- [3] V. Balan, M. Neagu, *Jet Single-Time Lagrange Geometry and Its Applications*, John Wiley & Sons, Inc., Hoboken, New Jersey, 2011.
- [4] I. Bucătaru, R. Miron, *Finsler-Lagrange Geometry. Applications to Dynamical Systems*, Romanian Academy Eds, Bucharest, 2007.
- [5] R. Miron, M. Anastasiei, *The Geometry of Lagrange spaces: Theory and Applications*, Kluwer Academic Publishers, 1994.
- [6] M. Neagu, C. Udrişte, *From PDEs systems and metrics to multi-time field theories and geometric dynamics*, Seminarul de Mecanică **79** (2001), Timișoara, Romania, 1-33.
- [7] C. Udrişte, *Geometric Dynamics*, Kluwer Academic Publishers, 2000.

Mircea Neagu
Department of Mathematics and Informatics
Transilvania University of Braşov
Blvd. Iuliu Maniu, No. 50, Braşov 500091, Romania
email: *mircea.neagu@unitbv.ro*